

The Condition Factor, Length-Weight Relationship and Abundance of *Elops senegalensis* (Regan, 1909) from Nkoro River, Niger Delta, Nigeria

J.F.N. Abowei

Department of Biological Sciences, Faculty of Science, Niger Delta University, Wilberforce Island, Amassoma, Bayelsa State, Nigeria

Abstract: The condition factor, length-weight relationship and abundance of *Elops senegalensis* from Nkoro River in the Niger Delta area of Nigeria was studied for a period of one year (January-December 2008). From a sample size of 1325 specimens, K value was 0.941 and the exponential equation was $Wt = 0.0153 (TL)^{3.066}$, indicating an isometric growth pattern. The highest catch was recorded in December (1.62), followed by August (1.41), July (1.28), January (1.12), February (1.01) and March (0.64), November (0.60), September (0.59), October (0.14), April (0.12) and May (0.11). *E. Senegalensis* occurred all year round during the study. The highest condition factor value (1.00) was recorded in January and the lowest (0.40) in October. The highest catch per unit effort (3.42) was recorded in stations 1, followed by station 2 (3.11), station 4 (2.11) and station 3 (0.12). Although *E. Senegalensis* from Nkoro River occurred all year round, the specie was more abundant in the dry season months of December and January; and also in the heavy rains months of July and August.

Key words: Abundance, condition factor, *Elops senegalensis*, length-weight relationship, Nigeria and Nkoro River

INTRODUCTION

The Senegalese ladyfish, *Elops senegalensis*, is a species of fish in the Elopidae family, and is native to the coastal waters of the eastern Atlantic Ocean from Mauritania to the Democratic Republic of the Congo. It is often confused with the West African ladyfish, *Elops lacerta*, can be distinguished only by the number of gill rakers on the lower part of the first gill arc, and the number of scales on the lateral line. The Senegalese ladyfish grows to a maximum total length of 90 centimeters and a maximum weight of 5.9 kilograms. Elopidae is a family of ray-finned fish containing the single genus *Elops*. They are commonly known as lady fishes, skip jacks, Jack-Rashes, or ten pounders.

The lady fish are a coastal dwelling fish found throughout the tropical and sub-tropical regions. Spawning takes place at sea and the fish larvae migrate inland entering brackish waters. Their food is smaller fish and crustaceans (shrimp). Typically throughout the species, the maximum size is 1metre (3.3ft) and the maximum weight 10kilograms (22lb). The body is fusiform (tapering spindle shape) and oval in cross-section; being slightly laterally compressed, the eyes are large and partially covered with fatty (adipose) eyelids. Like those of eels, the larvae are leptocephalic - being highly compressed, ribbon-like and transparent. After initial growth they shrink and then metamorphoses into the adult form. This family is fished, but the body is bony and therefore not edible by humans. However, the fish may be ground down for fish meal.

Condition factor compares the wellbeing of a fish and is based on the hypothesis that heavier fish of a given length are in better condition (Bagenal and Tesch, 1978). Condition factor has been used as an index of growth and feeding intensity (Fagade, 1979). Condition factor decrease with increase in length (Bakare, 1970; Fagade, 1979); and also influences the reproductive cycle in fish (Welcome, 1979). Condition factors of different species of cichlid fishes have been reported by Siddique (1977), Fagade (1978, 1979, 1983), Dadzie and Wangila (1980), Arawomo (1992) and Oni *et al.* (1983). Some condition factors reported for other fish species include; Alfred-Ockiya (2000), *Chana chana* in fresh water swamps of Niger Delta and Hart (1997), *Mugil cephalus* in Bonny estuary, Hart and Abowei (2007), ten fish species from the lower Nun River and Abowei and Davies (2009), *Clarotes lateiceps* from the fresh water reaches of the lower Nun river.

The length-weight relationship of fish is an important fishery management tool. Its importance is pronounced in estimating the average weight at a given length group (Beyer, 1987) and in assessing the relative well being of a fish population (Bolger and Connoly, 1989). Consequently, length - weight studies on fish are extensive. Notable among these are the reports Shenouda *et al.* (1994), for *Chrysichthys* spp. from the Southernmost part of the River Nile (Egypt), Alfred-Ockiya and Njoku (1995) for mullet in New Calabar River, Ahmed and Saha (1996) for carps in Lake Kapital, Bangladash, King (1996) for Nigeria fresh water fishes, Hart (1997) for *Mugil cephalus* in Bonny Estuary; Diri (2002) *Tilapia guineensis* in Elechi creek.

Catch Per Unit Effort (CPUE) is a useful index in the assessment of abundance of fish species (Gulland, 1975). It is essential in the determination of Maximum Sustainable Yield (MSY) and potential yield. Tobor (1992) reported that the inshore waters of most parts of the West African coast are rich in fish resources in quantities that can support commercial exploitation on a sustainable basis. However, later developments in fisheries studies have pointed to the depletion of the fish stocks (Okpanefe, 1987).

Accurate fisheries statistics in the river; and its adjoining flood plains is vital for the formulation of a sound fisheries management programme in the Nkoro River and similar water bodies. But, this is completely lacking. A part from (Scott, 1966; Reed *et al.*, 1967; Otobo, 1981; FAO, 1994; Otobo, 1993; Ita and Medahili, 1997; Sikoki and Otobotekere, 1999; Ezekiel *et al* 2002; Abowei and Ezekiel, 2003; Abowei *et al.*, 2007; 2008; Abowei and Hart 2007, 2008; 2009; Abowei and Davies, 2009), different water bodies, there are no reliable data on the abundance of *Elops senegalensis* from Nkoro River. This is essential for formulation of development plan in the fishing industry. This research therefore provides information to fill that gap in Nkoro River fisheries.

Unfortunately, no work has been done on the length-weight relationship of *Elops senegalensis* from the Nkoro River. A study of the abundance, condition factor and length-weight relationship of *E. senegalensis* from the Nkoro River adds more information on the family, pristigasteridae to complement the existing data in the management and culture of the species in the Nkoro River, Niger Delta.

MATERIALS AND METHODS

Study Area: The Nkoro River is a distributory of the Andoni River in the Niger Delta area of Nigeria. The Nkoro River lies between latitudes 4°28' to 4°45' N and longitudes 7°45' E. The Niger Delta is one of the world largest wetlands covering an area of approximately 70,000 km². The area is economically important and rich in biodiversity. Numerous activities such as oil exploration and production and agricultural activities go on in the region. Most of Nigeria's oil and gas reserves and production, which account for over 80% federal government's revenue, is located within the Niger Delta region. The Red and white mangroves (*Rhizophora* and *Avicennia* spp.) mangrove swamps and flood plains border the river and its numerous creeks; and these are well exposed at low tides.

Fish Sampling: Fish specimens were procured from artisanal fishers and middlemen at their landing site for the study. Sampling of landed catches was done twice in a month for a period of twelve months. The fishers used a wide range of fishing gear such as hook and line, long line, cast nets, gill nets and traps. From the catches, fish specimen were randomly identified using keys and descriptions by Holden and Reed (1972), Loveque *et al.*

(1991) and Reed *et al.* (1967). Specimens were stored in coolers containing ice and transported to the laboratory for further analysis.

The condition factor (k) of the experimental fish was estimated from the relationship:

$$K = \frac{100 W}{L^3} \quad (1)$$

Where K= condition factor,
W= weight of fish
L= length of fish (cm).

The Total Length (TL) of the fish was measured from the tip of the anterior or part of the mouth to the caudal fin using meter rule calibrated in centimeters. Fish were measured to the nearest centimeter. Fish weight was measured after blot drying with a piece of clean hand towel. Weighing was done with a tabletop weighing balance, to the nearest gram. The length measurements were converted into length frequencies with constant class intervals of 2cm. The mean lengths and weights of the classes were used for data analysis, the format accepted by FISAT (Gayanilo and Pauly, 1997).

The relationship between the length (L) and weight (W) of fish was expressed by equation (Pauly, 1983):

$$W = aL^b \quad (2)$$

Where W=Weight of fish in(g)
L=Total Length (TL) of fish in(cm)
a=Constant (intercept)
b=The Length exponent (slope)

The "a" and "b" values were obtained from a linear regression of the length and weight of fish. The correlation (r²), that is the degree of association between the length and weight was computed from the linear regression analysis:

$$R = r^2 \quad (3)$$

Abundance was estimated from the weight (kg) of the total catch of each station for each species over the period of this study and compared for difference using Analysis of Variance (ANOVA) to test for difference between the stations. Catch per unit effort was calculated by dividing the total monthly catch by the effort (number of fishers per boat) and finally dividing by the number of hours of fishing giving:

$$\begin{aligned} \text{CPUE} &= \text{Total catch/No of fishers/fishing hours} \\ \text{CPUE} &= \text{Kg/man/hr (King, 1991).} \end{aligned}$$

The figures for catch per unit effort were tested for variation on monthly and station basis using ANOVA.

Table 1: Condition factor and exponential equation of *Elops senegalensis* in Nkoro River

N	K	Exponential Equation
1325	0.941	Wt = 0.0153(TL) ^{3.066}

Table 2: Monthly catch per unit effort for *Elops senegalensis* in Nkoro River

Jan	Feb.	Mar.	Apr.	May	Jun.	Jul.	Aug.	Sep.	Oct.	Nov.	Dec.
1.12	1.01	0.64	0.12	0.11	0.10	1.28	1.41	0.59	0.14	0.60	1.62

Table 3: Catch per unit effort of *Elops senegalensis* at each station in Nkoro River

Station 1	Station 2	Station 3	Station 4
3.42	3.11	0.12	2.11




Fig. 1: Monthly condition factor for *E. senegalensis*

RESULTS

Table 1 expresses the condition factor and the exponential equation from the length weight relationship of *Elops senegalensis* in Nkoro River. From a sample size of 1325 specimens, K value was 0.941 and the exponential equation was $Wt = 0.0153 (TL)^{3.066}$, indicating an isometric growth pattern.

The monthly catch per unit effort for *Elops senegalensis* in Nkoro River is presented in Table 2. The highest catch was recorded in December (1.62), followed by August (1.41), July (1.28), January (1.12), February (1.01) and March (0.64), November (0.60), September (0.59), October (0.14), April (0.12) and May (0.11). *E. Senegalensis* occurred all year round during the study.

Fig. 1 shows the monthly condition factor for in *E. Senegalensis* Nkoro River. The highest condition factor value (1.00) was recorded in January and the lowest (0.40) in October.

The catch per unit effort of *Elops senegalensis* at each station in Nkoro River is presented in Table 3. The highest catch per unit effort (3.42) was recorded in stations 1, followed by station 2 (3.11), station 4 (2.11) and station 3.

DISCUSSION

The mean condition factors 0.70 and monthly condition factor ranging from 0.4-1.00 obtained in this study varied slightly with the results from other studies. Ajayi (1982), reported $K = 0.77-0.81$ for *Claroetes filamentosus* in lake Oguta; Nwadiaro and Okorie (1985) obtained $K = 0.49-1.48$ in Andoni river. The value obtained from the study showed that all species studied were in good condition. Gayanilo and Pauly (1997) reported that certain factors often affect the well-being of a fish. These include: data pulling, sorting into classes, sex, stages of maturity and state of the stomach.

The condition factor (K) reflects, through its variations, information on the physiological state of the fish in relation to its welfare. From a nutritional point of view, there is the accumulation of fat and gonad development (LeCren, 1951). From a reproductive point of view, the highest K values are reached in some species (Angelescu *et al.*, 1958). K also gives information when comparing two populations living in certain feeding, density, climate, and other conditions; when determining the period of gonad maturation; and when following up the degree of feeding activity of a species to verify whether it is making good use of its feeding source (Weatherley, 1972). From the above assertions we could conclude that the five species in this work reproduce between May to October since they recorded the lowest K at about this period.

Furthermore, Vazzoler (1996) confirmed that lowest K values during the more developed gonad stages might mean resource transfer to the gonads during the reproductive period. Braga (1986), through other authors, showed that values of the condition factor vary according to seasons and are influenced by environmental conditions. The same may be occurring in the environment under study since the floodplain is influenced by many biotic and abiotic factors, which favor the equilibrium of all the species in the ecosystem.

The values obtained for the weight – length relationship ($Wt = 0.0153 (TL)^{3.066}$) showed that *Elops senegalensis* was isometric in growth. Several authors have reported both isometric and allometric growth for different fish species from various water bodies. King 1991, reported allometric growth patterns for Tilapia species from Umuoseriche lake. King (1996) reported isometric growth for *Pseudotolithus elongatus* from Qua Iboe estuary. Ekeng (1990) also reported an isometric growth pattern for *Etmalosa fimbriata* from Cross River estuary in Cross River state. Marcus (1984), obtained an isometric growth patterns for *E. fimbriata* from coastal and brackish water of Akwa Ibom state. Sheneuda *et al.* (1994) also observed an isometric growth patterns for *Chysichthys auratus* from the southern most parts of River Nile and Egypt.

The transformed length fitted over weight gave linear growth indicating the three dimensional growth structures of most fish species (Lagler *et al.*, 1977). Values of the length exponent in the length-weight relationship being isometric implies that the fish species did not increase in weight faster than the cube of their total lengths. However, the weight of the rest species increased faster than the cube of their total lengths.

Length-weight relationships give information on the condition and growth patterns of fish (Bagenal and Tesch, 1978). Fish are said to exhibit isometric growth when

length increases in equal proportions with body weight for constant specific gravity. The regression co-efficient for isometric growth is '3' and values greater or lesser than '3' indicate allometric growth (Gayanilo and Pauly, 1997).

The catch per unit effort data of December (1.62), August (1.41), July (1.28), January (1.12), February (1.01) and March (0.64), November (0.60), September (0.59), October (0.14), April (0.12) and May (0.11) obtained from this study; varied from the results obtained from other studies. Scott (1966) reported that, rivers, lakes and swamps of the Niger Delta produced about 2,000 tonnes of fish per year. Moses (1981) estimated a mean annual catch of 4,791 tonnes from the cross river over a period of twelve years. Sikoki and Hart (1999) in the Brass river, estimated the total biomass of 160.20 of fish per boat, total catch of 254,554kg, annual production of 610.93 tonnes, estimated mean catch per boat of 384.90kg and a standing stock of 1.19km².

Variation in the total estimate values Nkoro River could be attributed to differences in fishing and industrial activities in the different rivers. The reason for the low estimates in the Nkoro River could be as a result of high mortality of both juveniles and brood stock of various fish species as a result of predatory activities, which is typical of the study area. A similar remark was made by, Ssentengo *et al.* (1986). Satia (1990) also noted the controversy surrounding fish production statistics. In the lower Nun River, much of the problem hampering the acquisition of reasonably accurate fisheries statistics and resource appraisal appear to stem mainly from lack of, or inadequate investment and lack of trained personnel to handle data collection.

Factors affecting fish distribution and abundance have already been reported by different workers. Availability of food, spawning rates, breeding grounds coupled with shelter, presence of current, vegetation, depth of water, breeding habits, migration and low predation have been suggested as major limiting factors affecting the distribution and abundance of various fish families in Kainji Lake (Ita, 1978).

Angelescu *et al.* (1958) reported fish catch varied with type of gear used, tidal condition and period of capture, diurnally and seasonally. From the work of King (1991), it is clear that most commercially and scientifically important fish species occurring in the Niger Delta waters can be landed all year round by artisanal fishers but there are months when they are more abundant.

REFERENCES

Abowei, J.F.N. and E.N. Ezekiel, 2003. Aspects of Some Physicochemical Parameters Of The Fresh Water Reaches of the Lower Num River, Niger Delta. *Int. J. Sci. Technol.*, 1(2): 5-9.

- Abowei, J.F.N. and A.I. Hart, 2007. Size, Composition, age, growth, mortality and exploitation rate of *Chysichthys nigrodigitatus* from Nun River, Niger Delta, Nigeria. *Afr. J. Appl. Zool. Environ. Biol.*, (9): 44-50.
- Abowei, J.F.N., F.D. Sikoki, A.I. Hart and M.E. Allison, 2007. Catch composition and seasonality of Fin fish in the fresh water reaches of the lower Nun River, Niger Delta. *J. Field Aquat. Stud.*, 3: 8-21.
- Abowei, J.F.N., C.C. Tawari, A.I. Hart and D.U. Garicks, 2008. Finfish species composition, abundance and distribution in the lower Sombreiro River, Niger Delta, Nigeria. *Int. J. Trop. Agric. Food Syst.*, 2(1): 46-53.
- Abowei, J.F.N. and A.I. Hart, 2008. Artisanal fisheries characteristics of the fresh water reaches of lower Nun River, Niger Delta, Nigeria. *J. Appl. Sci. Environ. Manage.*, 12(1): 5-7.
- Abowei, J.F.N. and A.I. Hart, 2009. Some morphometric parameters of ten finfish species from the lower Nun River, Niger Delta, Nigeria. *Res. J. Biol. Sci.*, 4(3): 282-288.
- Abowei, J.F.N. and A.O. Davies, 2009. Some population parameters of *Clarotes laticeps* (Rupell, 1829) from the fresh water reaches of the lower river, Niger Delta, Nigeria. *Am. J. Sci. Res.*, (2): 15-19.
- Ahmed, K.K. and S.B. Saha, 1996. Length-weight relationship of major carps in Kaptai lake. *Bangladesh. NAGA. The ICLARM Q.*, 19(2): 28.
- Angelescu, V., F.S. Gneri and A. Nani, 1958. *La merluza del mar argentino (biologia e taxonomia)*. *Secr. Mar. Serv. Hidrog. Nav. Publico*, H1004: 1-224.
- Ajayi, T., 1982. The age and growth of the tongue sole, *Cynolossus canariensis* (Stend, 1982). In: *Proceedings of the 2nd Annual Conference of the Fisheries Society of Nigeria (FISON) New Bush source*. 2.19.
- Alfred-Ockiya, J.F. and D.C. Njoku, 1995. A comparative analysis of the length weight relationship and condition factors of four species of grey mullet (pisces/mugilidae) from New Calabar River Rivers State, Nigeria. *J. Tech. Educ.*, pp: 5-10.
- Alfred-Ockiya, J.F., 2000. The length-weight relationship of snake head (*Chana chana*) from the fresh water swamps of Niger Delta. *J. Aquat. Sci.*, 15: 12-14.
- Arawomo, G.A.A., 1982. The growth of *Sarotherodon niloticus*. In: *Proceedings of The 2nd Annual Conference of the Institute*. New Bussa, Nigeria, pp: 221-227.
- Bagenal, T.B. and A.T. Tesch., 1978. *Conditions and Growth Patterns in Fresh Water Habitats*. Blackwell Scientific Publications, Oxford.
- Bakare, O., 1970. Bottom Deposits as Food of Inland Fresh Water Fish. In: *Kainji, A Nigerian Man-Made Lake*. S.A. Visser, (Ed.), *Kanyi Lake Studies Vol. 1. Ecology* Published for the Nigerian Institute.

- Beyer, J.E., 1987. On length-weight relationship computing the mean weight of the fish of a given length class. *Fishbyte*, 5(1): 11-13.
- Bolger, T. and P.L. Connolly, 1989. The selection indices for the measurement and analysis of fish condition. *J. Fish Biol.*, 17(3): 1-182.
- Braga, F.M.S., 1986. Estudo entre o factor de condição e relação peso/comprimento para alguns peixes marinhos. *Rev. Brasil. Biol.*, 46(2): 339-346.
- Diri, M.S., 2002. Length-weight relationship of *Sarotheredon melanotheron* and *Tilapia guineensis* in Elechi creek Niger Delta, Nigeria B.Sc. Project Rivers State University of Science and Technology, Port Harcourt, pp: 33.
- Dodzie, S. and B.C.C. Wangila, 1980. Reproductive biology, length-weight relationship and relative condition of pond raised tilapia zilli (Gervas). *J. fish Biol.*, 17: 243-253.
- Ekeng, E.O., 1990: Length-weight and diet composition of *Ethinalose fimbriata* (Bowchch). *Pliscea: clupeidae* in Cross River estuary, Nigeria. B.Sc. Project University of Calabar. Cross River State, Nigeria, pp: 36.
- Ezekiel, E.N., J.F.N. Abowei and A.I. Hart, 2002. The fish species assemblage of Odhioku-Ekpeye flood plains, Niger Delta. *Int. J. Sci. Technol.*, 1(1): 54-59.
- Fagade, S.O., 1978. Age determination of *Tilapia melanotheron* (Ruppel) in the Lagos lagoon, Nigeria. *International Symposium on Ageing of Fish* (In Bagenal), Tesch, pp: 71-77.
- Fagade, S.O., 1979. Observation of the biology of two species of *Tilapia* from the Lagos lagoon Nigeria. *Bull. Inst. Fond Afr. Nore* (Ser. A) 41: 627-658.
- Fagade, S., 1983. The biology of *chromido Tilapia guntheri* from a small lake. *Arch. Hydobil.*, 97: 60-72.
- Food and Agricultural Organization (FAO), 1994. Report of the study on exploration and use of small *Pelagic* spp. in West Africa, FAO Fish Circular No. 880, pp: 18-28.
- Gayaniilo, F.C. and D. Pauly, 1997. FAO ICLARM stock assessment tools (FISAT): References Manual, FAO Computerized Information Series (Fisheries) (8): 262.
- Gulland, J.A., 1975. Manual of Sampling and statistical methods for fisheries biology. Part 1, Sampling Methods. FAO Manuals in Fisheries Science. No. 3 FRS/M 3: 1-87.
- Hart, S.A., 1997. The Biology of *Mugil cephalus* in Bonny River estuary. M.Sc. Thesis University of Port Harcourt, Nigeria, pp: 42.
- Hart, A.I. and J.F.N. Abowei, 2007. A study of the length-weight relationship, condition factor and age of ten fish species from the lower Nun river. *Niger Delta. Afr. J. Appl. Zool. Environ. Biol.*, 9: 13-19.
- Holden, M. and W. Reed, 1972. *West African Fresh Water Fishes*. Longmans Ltd., London, pp: 33.
- Ita, E.O., 1978. An analysis of fish distribution in Kainji Lake, Nigeria. *Hydrobiologia*, 58: 233-244.
- Ita, E.O and I. Maelahili, 1997. The current status of fish stock and fisheries in Kainji Lake; consultancy report on fish stock assessment in Kainji Lake. The Nigerian-German (GTZ) Kainji Lake Fisheries Promotion Project, New Bussa, pp: 128.
- King, R.P., 1991. The biology of *tilapia mariae* Bovlenger 1899 (Perciformes: Cicchliidae) in a Nigeria Rainforest stream. Ph.D. Thesis, Department of Zoology, University of Port Harcourt, Nigeria.
- King, R.P., 1996. Population dynamics of the mud skipper *Periophthalmus barbarus* (Gobidae) in the estuarine swamps of Cross River Nigeria. *J. Aquat. Sci.* 11: 31-34.
- Lagler, K.F., J.E. Bardach, R.R. litter and D.R.M. Passimo, 1977. *Ichthyology*. John Wiley and Sons Inc., pp: 506.
- LeCren, E.D., 1951. The length-weight relationship and seasonal cycle in gonad weight and condition in the perch *Perca fluviatilis*. *J. Anim. Ecol.*, 20(2): 201-219.
- Loveque, C., O. Pyugy and G.G. Teugels, 1991. The fresh and brackish Water fishes of West Africa. *Musee Royale de l'Afrique Centrale*. Tervurem, Belgique, Editions de l'ORSTOM, 1: 384.
- Marcus, C., 1984. Biology of bonga fish; *Ethmalosa fimbriata* (Bowdich) in the Nigeria Coastal and Brackish waters project (NF) 1,2. Annual Report, Nigeria Institute for Oceanography and Marine Research Lagos, pp: 232.
- Moses, B.S., 1981. Preliminary estimates of potential yield of Cross River State inland fresh waster fisheries. In: *Pro. First CR. Fish Cont. MFN Calabar*, pp: 41-46.
- Nwadiaro, C.S. and P.U. Okorie, 1985. Biometric characteristics: length weight relationships and condition factors in *Chrychthys filamentosus*, Pisces, Bagandae from Oguta lake Nigeria. *Biol. Afr.*, 2: 48-56.
- Okpanefe, M.O., 1977. Agriculture and National Resources Fisheries Statistics survey of Nigeria. *Fed. Rep. of Nig. Occassional Paper*, 23: 5.
- Oni, S.K., J.Y. Olayemi and J.D. Adegboye, 1983. The comparative physiology of three ecologically (Rupel). *Synodonts schall*. Block and Schneider and *Tilapia zilli* (Gervais). *J. Fish. Biol.*, 22: 105-109.
- Otobo, A.J.T., 1981. Identification of fish species in a stretch of River Nun. HND Project Rivers State University of Science and Technology Port Harcourt.
- Otobo, A.J.T., 1993. The ecology and fishery of the Pygmy Herring. *Sierrathensa Leonensis* (Thysvan Den Audenaerde, (1969) in the Nun River and Taylor Creek of the Niger Delta. Ph.D. Thesis. University of Port Harcourt, pp: 298.
- Pauly, D., 1983. Some simple methods for the assessment of tropical fish stock. *FAO Fish. Tech. Paper No.* 234, pp: 52.

- Reed, W., T.A.J. Burchad, J. Hopson, J. Jenness and I. Yaro, 1967. Fish and fisheries of Northern Nigeria. Ministry of Agriculture, Northern Nigeria, pp: 226.
- Satia, B.P., 1990. National review for aquaculture development in Africa. Nigeria. FAO Fish Girel: 770(29): 191.
- Scott, J.S., 1966. Report on the Fisheries of the Niger Delta Special Area. NDDB, pp: 109.
- Shenouda, T.S., F.A. Faten, M.R. Mahmoud and M.M. Ray, 1994. A detail study on age and growth for *Chrysichthys auratus* and *Chrysichthys rueppelli* from the southern most part of the River Nile (Egypt). J. Egypt Ger. Soc., 200(1412): 73-101.
- Siddique, A.Q., 1977. Reproductive biology, length-weight and relative condition of *Tilapia leucostica* (Trewaeva in lake Naivasha, Kenya). J. Fish. Biol., 10: 351-260.
- Sikoki, F.D. and S.A. Hart, 1999. Studies on the fish and fisheries of the brass river system and adjoining coastal waters in Bayelsa State, Nigeria. J. Appl. Sci. Environ. Manage., 2: 63-67.
- Sikoki, F.D. and A.J.T. Otobotekere, 1999. Fisheries. In: The Land People of Bayelsa State Central Niger Delta. Alagoa E.C., (Ed.). Port Harcourt, pp: 301-319.
- Ssentengo, G.W., E.T. Ukpe and T.O. Ajayi, 1986. Marine fishery resources of Nigeria. Reviews of exploited fish stock CECAF/ECAC series, 86/40/FAO. Rome, pp: 52.
- Tobor, J.G., 1992. Fin and shell fish of conservation interest in Nigeria. NIOMR Tech Paper No. 79, pp: 1-23.
- Vazzoler, A.E., 1996. Biologia da reprodução de peixes Teleosteos: teoria e prática. EDUEM, SBI, Maringá, pp: 169.
- Weatherley, A.H., 1972. Growth and Ecology of Fish Populations. Academic Press, London, pp: 293.
- Welcome, R.L., 1979. Fisheries Ecology of Flood Plain Rivers. Longman Press, London, pp: 317.